

Instructions for use HUMIN PLUS for Corn



Instructions for use of HUMIN PLUS when processing corn.

Application HUMIN PLUS:

- Microfertilizer HUMIN PLUS is used as an aqueous solution for the processing of crops alone or in conjunction with other fertilizers and crop protection agents.

INSTRUCTION on the use of "HUMIN PLUS" in soy processing

Treatment of planting material:

- Consumption of products - 200 ml per 10 liters of water. Planting material is processed with Humin Plus in any convenient way: soaking, spraying, pickling tank mixing, etc., and after treatment planting seeds to stand at least 1 hour.

Solution preparation:

- The first treatment for vegetation - 400 ml per 1 ha
- The second treatment on vegetation - 300 ml per 1 ha.

Method of application:

- Humin Plus is added to the spray tank. The optimum concentration of the working solution for processing crops - 1:1000. Concentration may vary depending on the settings of the sprayer but should be based on the recommended application rates per hectare.

Seed treatment is obligatory.

Plant Growth:

- During the onset of the main phases of plant development:
- The first treatment for the growing season – 3 to 5 leaves.
- The second treatment on vegetation - during budding. The best treatment time from 5 to 10 am and then from 16 to 21 o'clock in the evening. The second treatment - not earlier than 14 days after the first. The working solution should be prepared immediately before use.